

Mark Scheme (Results)

October 2021

Pearson Edexcel International A Level In Decision Mathematics (WDM11) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <u>www.edexcel.com</u> or <u>www.btec.co.uk</u>. Alternatively, you can get in touch with us using the details on our contact us page at <u>www.edexcel.com/contactus</u>.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2021 Question Paper Log Number P69351A Publications Code WDM11_01_2110_MS All the material in this publication is copyright © Pearson Education Ltd 2021

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

- 1. The total number of marks for the paper is 75.
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
- M marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
- A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
- **B** marks are unconditional accuracy marks (independent of M marks)
- Marks should not be subdivided.
- 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod benefit of doubt
- ft follow through
- the symbol $\sqrt{\text{ will be used for correct ft}}$
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- dep dependent
- indep independent
- dp decimal places
- sf significant figures
- ***** The answer is printed on the paper
- The second mark is dependent on gaining the first mark
- 4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
- 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
- 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
- 7. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme	Marks
1.(a)	A path is a (i) finite sequence of edges, such that (ii) the end vertex of one edge in the sequence is the start vertex of the next, and in which (iii) no vertex appears more than once	B2, 1, 0 (2)
(b)	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	M1 A1 (ACBFD) A1 (GE) A1ft (HJ)
	Shortest path: A B E H J Length: 33 (km)	Al Alft (6)
(c)	Shortest path from J to A via G: J G D C A	B1
	Length: $20 + 15 = 35$ (km)	B1ft (2)
		10 marks

Number	Scheme	Marks
	Notes for Question 1	

a1B1: One of the three points made clearly ('finite, edges', 'end vertex of one edge is the start vertex of the next', 'no vertex appears more than once')

a2B1: All three points made clearly. Candidates who state that a path is a walk in which no vertex appears more than once can score B1B0 only

In (b) it is important that all values at each node are checked very carefully – the order of the working values must be correct for the corresponding A mark to be awarded e.g. at F the working values must be 15 14 13 in that order (so 15 13 14 is incorrect). It is also important that the order of labelling is checked carefully. The order of labelling must be a strictly increasing sequence – so 1, 2, 3, 3, 4, ... will be penalised once (see notes below) but 1, 2, 3, 5, 6, ... is fine. Errors in the final values and working values are penalised before errors in the order of labelling

b1M1: Working values - a larger value replaced by a smaller value for at least two of the five activities D, E, F, G, J

b1A1: All values at A, C, B, F and D correct and the working values in the correct order

b2A1: All values at G and E correct and the working values in the correct order

b3A1ft: All values in H and J correct on the follow through and the working values in the correct order **b4A1:** cao (A B E H J only)

b5A1ft: Follow through on their final value at J **only** (condone lack of units)

c1B1: cao (J G D C A only)

c2B1ft: 35 or follow through their final value at G + 15

Question Number	Question Scheme						
2.	$2. \qquad y \ge 3x$						
	$z - x \ge 50$	B1					
	<i>y</i> ≤120	B1					
	Sub. $x + y + z = 180$	M1					
	$2x + y \le 130$						
	Maximise $(P=)x+y$	B1 (6)				
1B1: cao ($(y \ge 3x)$ oe (two terms only with integer coefficients)						
2B1: cao $(z - x \ge 50)$ – may be implied by later working oe (three terms only with integer coefficients) 3B1: $y \le 120$ oe							
1M1: Eliminating z by substituting $x + y + z = 180$ into an inequality that involves z and x only							
1A1: $2x + y \le 130$ oe (three terms only with integer coefficients)							
4B1: correct objective with 'maximise' or 'max' but not 'maximum' – either the expression $x + y$ or any other letter for <i>P</i> except <i>x</i> , <i>y</i> or <i>z</i>							

Question Number	Scheme	Marks					
3. (a)	Prim: AE, EG, CE; DG, CF; DH, BF	M1 A1 A	1 (3)				
(b)	Weight of MST = 197	B1	(1)				
(c)	Initial upper bound = $2(197) = 394$	B1ft	(1)				
(d)	A - E - G - D - H - B - F - C - A	M1					
(u)	23 + 24 + 26 + 33 + 38 + 34 + 32 + 38 = 248	A1					
	A-E-G-D-H-F-C-B-A	A1					
	23 + 24 + 26 + 33 + 38 + 32 + 35 + 36 = 247	A1	(4)				
(e)	247	B1ft	(1)				
(f)	Weight of RMST is 174	B1ft					
	Lower bound = $174 + 23 + 35 = 232$	M1 A1	(3)				
(g)	$232 \square$ optimal value $\square 247$	M1 A1	(2)				
	Notes for Question 3						

a1M1: Prim's – first three arcs correctly chosen in order (AE, EG, CE, ...) **or** first four nodes {A, E, G, C, ...} correctly chosen in order. If any explicit rejections seen at some point then M1 (max) only. Order of nodes may be seen at the top of a matrix/table {1, -, 4, -, 2, -, 3, -}. Starting at any other node can score M1 only for first three arcs chosen correctly

a1A1: First five arcs correctly chosen in order (AE, EG, CE, DG, CF, ...) **or** all eight nodes {A, E, G, C, D, F, H, B} correctly chosen in order. Order of nodes may be seen at the top of a matrix so for the first two marks accept {1, 8, 4, 5, 2, 6, 3, 7} (**no** missing numbers)

a2A1: cso – all **arcs** correctly **stated** and chosen in the correct order (with no additional arcs). They must be considering arcs for this final mark (do not accept a list of nodes or numbers across the top of the matrix unless the correct list of arcs (in the correct order) is also seen)

b1B1: cao (197 – ignore units) should come from 23 + 24 + 25 + 26 + 32 + 33 + 34**c1P1ft:** Follow through double their ensurer to (b)

c1B1ft: Follow through double their answer to (b)

Mark (d) and (e) together

d1M1: Nearest neighbour starting at A with first five nodes correct (A - E - G - D - H -)

d1A1: One correct route (must return to A)

d2A1: One correct value or both correct routes

d3A1: Both correct values (do not isw if values doubled) and both correct routes (must both return to A)

SC in (d) correct Hamiltonian paths and corresponding weights (AEGDHBFC (210) and AEGDHFCB (211)) scores M1A1A0A0

e1B1ft: Follow through their least weight route from (d) – must have or imply two Hamiltonian cycles in (d) or (e)

f1B1ft: Either 174 or 24 + 25 + 26 + 32 + 33 + 34 or 197 - 23 or the weight of their MST from (b) - 23 **f1M1:** Weight of RMST + 23 + 35 (two smallest arcs incident to A) with 151£ RMST£ 197 (if clearly not six arcs in RMST then M0)

f1A1: cao (232) – if correct answer with no working then awarded B0M1A1 – as a minimum for full marks accept 174 + 23 + 35 = 232 but 174 + 58 = 232 scores B1M1A0

g1M1: Any indication of an interval from their answer to (f) to their answer to (e) with one value correct g1A1: cao (either 232 £ optimal value £ 247 or 232 < optimal value £ 247)

Question Number	Scheme	Marks						
4. (a)	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	B2, 1, 0 (2)						
(b)	v = 7, w = 4, x = 6, y = 16, z = 19	B3, 2, 1, 0 (3)						
(c)	$\frac{74}{25} = 2.96$ so 3 workers	B1 (1)						
(d)	e.g. 0 2 4 6 8 10 12 14 16 18 20 22 24 26 B E H L A C F J M D G K I	M1 A1 A1 (3)						
(e)	New minimum project completion time: 27 (days)	B1						
	Critical path: ACFK	B1 (2)						
	Notes for Question 4							
a1B1: Ang a2B1: cao	y 6 rows correct (not including A and B) (condone blank rows for A and B)							
b1B1: Any 2 correct valuesb2B1: Any 3 correct valuesb3B1: All 5 values correct								
c1B1: cao (3 from correct working) – as a minimum for correct working accept either 2.96 or $\frac{5+6+7+4+7+4+5+7+10+4+6+5+4}{25}$ or $\frac{74}{25}$ d1M1: Not a cascade chart. 4 workers used at most, at least 9 activities placed d1A1: 4 workers. All 13 activities present (just once). Condone two errors either precedence or time interval or activity length. An activity can give rise to at most three errors; one on duration, one on time interval and only one on IPA d2A1: 4 workers. All 13 activities present (just once). No errors e1B1: cao (27 only) e2B1: cao (ACFK or KFCA only)								

Question Number			Scheme	 Marks
Activit	y Duration	Time	IPA	
А	5	0-6	-	
В	6	0-6	-	
С	7	5 - 13	Α	
D	4	5-13	A	
E	7	6 – 13	A, B	
F	4	12 – 19	С	
G	5	12 - 20	С	
Н	7	13 - 20	C, D, E	
Ι	10	6 - 20	A, B	
J	4	6-21	A, B	
K	6	16 - 25	F	
L	5	20 - 25	F, G, H, I	
М	4	20 - 25	F, G, H, I, J	

Question Number	Scheme	Marks	
5.(a)	A(DG)C + D(GH)E = 12 + 9 = 21	M1 A1	
	AD + C(GH)E = 5 + 10 = 15*	A1	
	A(DGH)E + C(G)D = 14 + 7 = 21	A1	
	Repeated arcs: AD, CG, GH, EH	A1	
	Length of route: $166 + 15 = 181$ (km)	Alft	(6)
(b)	Vertex C: 3 times	B1	(1)
(c)	CD (7) is the shortest path between two odd nodes excluding A	M1	
	Repeat CGD (7) since this is the shortest path excluding A The route finishes at E	A1	
	Length of route = $166 + 7 = 173$ (km)	A1 ((3)
		10 marks	
	Notes for Question 5		

a1M1: Three distinct pairings of the correct four odd nodes (A, C, D, E)

a1A1: One row correct including pairings and totals

a2A1: Two rows correct including pairings and totals

a3A1: All three rows correct including pairings and totals

a4A1: The smallest repeat **arcs** (accept AD, CG, GH, EH only)

a5A1ft: Correct answer of 181 or 166 + their least

b1B1: cao (3)

c1M1: Identifies the need to repeat one path of the three (DE, CE, CD) which does not include A (this maybe implicit) **or** listing of only these three possible repeats. This mark is dependent on either scoring the M mark in (a) or stating all three possible paths in this part. As a minimum accept the stating of one of these three paths

c1A1: Identifies C(G)D as the least **and** E as the finishing point. They have to <u>explicitly state</u> that C(G)D is the <u>least</u> path of those that <u>do not include A</u> (this can be done by stating that CD is the least of CD, CE, DE only (so with no others) **or** stating that CD is the least of those that don't include A but not for just 'CD is the least')

c2A1: cao (173)



Notes for Question 6The lines in (a) must define the correct FR and pass within half a square of the points stated: $4x + 3y = 300$ with points (0, 100) and (25, 0) $x + y = 100$ with points (0, 0) and (60, 20) $xH + y = 100$ with points (0, 0, 0) and (60, 20) $xH = 2y = 130$ with points (0, 0, 0) and (60, 20) $xH = 2y = 130$ with points (0, 0, 0) and (60, 20) $xH = 2y = 130$ with points (0, 0, 0) and (60, 20) $xH = 2y = 130$ with points (0, 0, 0) and (60, 20) $xH = 2y = 130$ with points (0, 0, 0) and (60, 20) $xH = 2y = 130$ with points (0, 100) and (25, 0) $x = 2y = 130$ with points (0, 100) and (25, 0) $x = 2y = 130$ with points (0, 100) and (25, 0) $x = 2y = 130$ with points (0, 100) and (25, 0) $x = 2y = 130$ with points (0, 100) and (10, 20) $x = 2y = 130$ $x = 10 = 12y = 100 = 12y = 12y$	Question				Scheme	Marks						
The lines in (a) must define the correct FR and pass within half a square of the points stated: 4x + 3y = 300 with points (0, 100) and (75, 0) 4x + y = 130 with points (0, 65) and (130, 0) 3y = x with points (0, 65) and (60, 20) a IB : Any two lines correctly drawn a 3B : Any three lines correctly drawn a 3B : Any three lines correctly drawn a 3B : A correct objective line drawn on the graph with a gradient of $0.8 -$ intersections points with each axes given below $\frac{x + y}{10} = 10$ $\frac{x}{10} + \frac{y}{10}$ $\frac{x}{10} + \frac{y}{10} + \frac{10}{10}$ $\frac{x}{10} + \frac{y}{10} + \frac{10}{10}$ $\frac{x}{10} + \frac{y}{10} + \frac{10}{10} + 1$			Notes for Question 6									
4x + 3y = 300 with points (0, 100) and (75, 0) $4x + y = 100 with points (0, 100) and (25, 0)$ $x + 2y = 130 with points (0, 0) and (60, 20)$ a IB1 : Any two lines correctly drawn a 2B1 : Any two lines correctly drawn a 2B1 : Any two lines correctly drawn a 2B1 : All four lines correctly drawn a 4B1 : Correct <i>R</i> labelled – dependent on all three previous B marks bil B1 : A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each a a bit : A line intersection in the second se	The lines in (a) must define the correct FR and pass within half a square of the points stated:											
$\begin{aligned} 4x + y &= 100 \text{ with points } (0, 100) \text{ and } (25, 0) \\ x + 2y &= 130 \text{ with points } (0, 65) \text{ and } (130, 0) \\ 3y &= x \text{ with points } (0, 0) \text{ and } (60, 20) \\ all B1: Any three lines correctly drawn \\ a2B1: Any three lines correctly drawn \\ a3B1: All four lines correctly drawn \\ a3B1: Correct R labelled – dependent on all three previous B marks \\ bilB1: A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each axes given below\frac{x}{100} \frac{y}{12} \frac{y}{12.5} \frac{10}{100} \frac{125}{50} \frac{100}{100} \frac{125}{50} \frac{100}{125} \frac{100}{100} \frac{112.5}{50} \frac{100}{125} \frac{100}{1$	4x + 3y = 300 with points (0, 100) and (75, 0)											
$x + 2y = 130 \text{ with points } (0, 65) \text{ and } (130, 0)$ $3y = x \text{ with points } (0, 0) \text{ and } (60, 20)$ a BB : Any two lines correctly drawn a 3B : Any two lines correctly drawn a 3B : Any two lines correctly drawn a 3B : Correct <i>B</i> labelled – dependent on all three previous B marks b BB : A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each axes given below $\frac{x}{10} + \frac{y}{10}$ $\frac{x}{10} + \frac{y}{10} + \frac{x}{10}$ $\frac{x}{10} + \frac{y}{10}$ $\frac{x}{10} + \frac{x}{10} + \frac{x}{10}$ $\frac{x}{10} + \frac{x}{10} + \frac{x}{$	4x + y = 1	4x + y = 100 with points (0, 100) and (25, 0)										
$\begin{array}{llllllllllllllllllllllllllllllllllll$	x + 2y = 1	130 with point	ts (0, 65) a	nd (130, 0))							
al B1: Any three lines correctly drawn a2B1: Any three lines correctly drawn a4B1: Correct <i>R</i> labelled – dependent on all three previous B marks bill 1: A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each axes given below $\frac{x}{10} \frac{y}{20} \frac{1}{10} \frac{x}{25} \frac{y}{20} \frac{1}{25} \frac{1}{20} \frac{1}{25} \frac{1}{2} \frac{1}{$	3y = x w	ith points (0, 0)) and (60,	20)								
a2B1: Any three lines correctly drawn a3B1: All four lines correctly drawn a4B1: Correct <i>R</i> labelled – dependent on all three previous B marks bi1B1: A correct objective line drawn on the graph with a gradient of −0.8 – intersections points with each axes given below $\frac{x}{10} \frac{y}{10}$ $\frac{x}{10} \frac{y}{12.5}$ $\frac{x}{10} \frac{y}{2.5}$ $\frac{x}{20}$ $\frac{x}{10} \frac{y}{10}$ $\frac{x}{20} \frac{y}{16}$ $\frac{x}{12.5} \frac{y}{10}$ $\frac{x}{12.5} \frac{y}{12.5}$ $\frac{x}{12.5} \frac{x}{12.5}$ $\frac{x}{12.5} $	a1B1: Ang	y two lines con	rrectly drav	wn								
a3B1 : All four lines correctly drawn a4B1 : Correct <i>R</i> labelled – dependent on all three previous B marks biB1 : A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each axes given below a $\frac{x}{y}$ b $\frac{y}{10}$ a $\frac{x}{24}$ b $\frac{x}{12.5}$ b $\frac{y}{12.5}$ b $\frac{1}{25}$ c $\frac{1}{20}$ a $\frac{30}{24}$ b $\frac{1}{25}$ c $\frac{1}{20}$ a $\frac{1}{30}$ c $\frac{24}{40}$ b $\frac{1}{25}$ c $\frac{1}{20}$ b $\frac{1}{25}$ c $\frac{1}{20}$ c $\frac{1}{62.5}$ c $\frac{50}{60}$ b $\frac{60}{60}$ 48 b $\frac{75}{75}$ 60 b $\frac{60}{125}$ b $\frac{100}{125}$ b	a2B1: An	y three lines co	orrectly dra	awn								
adBi: Correct R labelled – dependent on all three previous B marks bilB1: A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each accs given below $ \frac{x}{10} + \frac{y}{10} + \frac{x}{12.5} + \frac{y}{10} + \frac{x}{10.5} + \frac{x}$	a3B1: All	four lines cor	rectly draw	vn								
bil B1: A correct objective line drawn on the graph with a gradient of -0.8 – intersections points with each axes given below $\frac{x y}{10 8}$ $\frac{x y}{10 8}$ $\frac{x y}{12.5 10}$ $\frac{25 20}{25 20}$ $\frac{30 24}{30 24}$ $\frac{37.5 30}{50 40}$ $\frac{50 40}{62.5 50}$ $\frac{60 48}{75 50}$ $\frac{75 60}{100 80}$ $\frac{90 72}{12.5 100}$ $\frac{112.5 90}{125 100}$ $\frac{112.5 90}{15 100}$ $\frac{112.5 90}{15 100 100 100 100 10$	a4B1: Con	rrect R labelle	d – depend	ent on all t	hree previous B marks							
x y 10 8 20 16 20 16 20 16 20 16 20 16 20 16 30 24 50 40 50 40 50 40 50 40 62.5 50 60 48 75 60 70 56 80 64 90 72 112.5 90 100 80 90 72 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112.5 90 112	bi1B1: A	correct objecti	ive line dra	wn on the	graph with a gradient of -0.8 – intersections point	nts with each						
$\begin{vmatrix} x & y \\ 10 & 8 \\ 20 & 16 \\ 30 & 24 \\ 40 & 32 \\ \hline 50 & 40 \\ 60 & 48 \\ \hline 75 & 50 \\ \hline 40 \\ \hline 50 & 40 \\ \hline 60 & 48 \\ \hline 75 & 50 \\ \hline 70 & 56 \\ \hline 80 & 64 \\ \hline 90 & 72 \\ \hline 100 & 80 \\ \hline 125 & 100 \\ \hline 100 & 80 \\ \hline 125 & 100 \\ \hline 100 & 80 \\ \hline 125 & 100 \\ \hline 100 & 80 \\ \hline 125 & 100 \\ \hline 112.5 & 90 \\ \hline 100 & 80 \\ \hline 125 & 100 \\ \hline 112.5 & 100 \\ \hline 113.1 & can (42, 44) only) - dependent on first three B marks in (a) and the first B mark in (b) \\ \hline 113.1 & can (42, 44) only) - dependent on first three B marks in (a) and the first B mark in (b) \\ \hline 113.1 & can (42, 44) only) - dependent on first three B marks in (a) and the first B mark in (b) \\ \hline 113.1 & can (77.6 only) - dependent on first three B marks in (a) and the first B mark in (b) \\ \hline 112.4 & can (77.6 only) - dependent on the marks in (a) and/or (b) \\ c1M1: Point testing method: 10k + 60 l kx_1 + y_1 or 60k + 20 l kx_1 + y_1 or 42k + 44 l kx_1 + y_1 where l is any inequality sign or the equals sign and (x_1, y_1) is their numerical V or (42, 44). Objective line method: -\frac{1}{2} - k \text{ or } -\frac{4}{3} - k \text{ or } \frac{1}{3} - k \text{ where } l is any inequality or equals. Or one correct answer stated \\ c1A1: One correct answer (k < \frac{1}{2}, k \in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3}) - if no method or working (as shown above) then A0 c2M1dep: Point testing: 10k + 60 l kx_1 + y_1 and 60k + 20 l kx_1 + y_1 where l is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their V (but not ((10, 60) or (60, 20))) (so V must now be the intersection of the two lines 4x + 3y = 300 and x + 2y = 130). $	axes giver	below										
$\begin{bmatrix} 10 & 8 \\ 20 & 16 \\ 30 & 24 \\ \hline \\ 40 & 32 \\ \hline \\ 50 & 40 \\ \hline \\ 50 & 40 \\ \hline \\ \hline \\ 50 & 40 \\ \hline \\ \hline \\ \hline \\ 75 & 60 \\ \hline \\ \hline \\ \hline \\ 75 & 60 \\ \hline \\ $	<i>x</i>	<u>y</u>	<i>x</i>	<u>y</u>								
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	10	8	12.5	10								
3024403250406048705680649072112.590125100billM1: Solving correct pair of simultaneous equations for their V (or if not labelled then the vertex consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective linebillA1: cao (42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b)billA1: cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b)SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks in (a) and/or (b)c1M1: Point testing method: 10k + 60□ kx ₁ + y ₁ or 60k + 20□ kx ₁ + y ₁ or 42k + 44□ kx ₁ + y ₁ where □ is any inequality sign or the equals sign and (x_1, y_1) is their numerical V or (42, 44). Objective line method:- $\frac{1}{2}$ □ - k or - $\frac{4}{3}$ □ - k where □ is any inequality or equals. Or one correct answer statedc1A1: One correct answer ($k < \frac{1}{2}$, $k > \frac{4}{3}$, $k > \frac{4}{3}$) - if no method or working (as shown above) then A0c2M1dep: Point testing: 10k + 60□ kx ₁ + y ₁ and 60k + 20□ kx ₁ + y ₁ where □ is any inequality sign or the equals. Or both correct answers stated with no workingc2A1: Both correct answers only ($k < \frac{1}{2}$ or $k \in \frac{1}{2}$ and $k > \frac{4}{3}$ or $k^3 - \frac{4}{3}$ □ - k and - $\frac{4}{3}$ □ - k	20	16	25	20								
$\frac{40}{50} \frac{32}{40}$ $\frac{50}{60} \frac{40}{48}$ $\frac{50}{70} \frac{40}{56}$ $\frac{61}{75} \frac{50}{60}$ $\frac{87.5}{70} \frac{70}{100}$ $\frac{80}{72} \frac{64}{112.5} \frac{90}{100}$ $\frac{112.5}{100}$ $\frac{100}{80}$ $\frac{112.5}{125} \frac{100}{100}$ $\frac{100}{80}$ $\frac{100}{125} \frac{100}{125}$ $\frac{100}{12} \frac{100}{125}$ $\frac{100}{12} \frac{100}{12}$ $\frac{100}{12$	30	24	37.5	30								
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	40	32	50	40								
60487560705610080907210080125100125bi2B1: cao V labelled – dependent on first three B marks in (a) and the first B mark in (b)bi1M1: Solving correct pair of simultaneous equations for their V (or if not labelled then the vertex consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective linebi1A1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b)SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks in (a) and/or (b)c1M1: Point testing method:10k + 60 □ kx ₁ + y ₁ or 60k + 20 □ kx ₁ + y ₁ or 42k + 44 □ kx ₁ + y ₁ where □ is any inequality sign or the equals sign and (x ₁ , y ₁) is their numerical V or (42, 44). Objective line method:- $\frac{1}{2}$ □ - k or - $\frac{4}{3}$ □ - k or $\frac{1}{3}$ □ - k where □ is any inequality or equals. Or one correct answer statedc1A1: One correct answer (k < $\frac{1}{2}$, k £ $\frac{1}{2}$, k > $\frac{4}{3}$, k 3 $\frac{4}{3}$) - if no method or working (as shown above) then A0c2M1dep: Point testing:10k + 60 □ kx ₁ + y ₁ and 60k + 20 □ kx ₁ + y ₁ where □ is any inequality sign or the equals. Or both correct answers stated with no workingc2A1: Both correct answers only (k < $\frac{1}{2}$ or k£ $\frac{1}{2}$ and k > $\frac{4}{3}$ or k $\frac{3}{3}$) with working as shown above	50	40	62.5	50								
70 56 87.5 70 80 64 100 80 90 72 125 100 112.5 90 12.5 100 123 100 112.5 90 125 100 112.5 90 125 100 112.5 90 125 100 112.5 90 125 100 112.5 100 1112.5 90 12.5 100 1112.5 90 12.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 1112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 112.5 100 <tr< td=""><td>60</td><td>48</td><td>75</td><td>60</td><td></td><td></td></tr<>	60	48	75	60								
8064100809012.590125100biB1: cao V labelled – dependent on first three B marks in (a) and the first B mark in (b)bii1M1: Solving correct pair of simultaneous equations for their V (or if not labelled then the vertex consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective line bii1A1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b)bii2A1: cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b)SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks carned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b)c1M1: Point testing method:10k + 60 □ kx ₁ + y ₁ or 60k + 20 □ kx ₁ + y ₁ or 42k + 44 □ kx ₁ + y ₁ where □ is any inequality sign or the equals sign and (x ₁ , y ₁) is their numerical V or (42, 44). Objective line method: - $\frac{1}{2}$ □ - k or - $\frac{4}{3}$ □ - k or $\frac{1}{3}$ □ - k where □ is any inequality or equals. Or one correct answer statedc1A1: One correct answer:(k < $\frac{1}{2}$, $k > \frac{4}{3}$, $k^3 = \frac{4}{3}$) - if no method or working (as shown above) then A0 c2M1dep: Point testing: 10k + 60 □ kx ₁ + y ₁ and 60k + 20 □ kx ₁ + y ₁ where □ is any inequality sign or the equals sign and (x ₁ , y ₁) is (42, 44) or their V (but not ((10, 60) or (60, 20)) (so V must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}$ □ - k and $-\frac{4}{3}$ □ - k where □ is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only ($k < \frac{1}{2}$	/0	56	8/.5	/0								
90 1/2 100 80 125 100 bill 1 : Solving correct pair of simultaneous equations for their <i>V</i> (or if not labelled then the vertex consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective line bill 1 : cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bill 1 : cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bill 2 : cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b) SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1 : Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). Objective line method: $-\frac{1}{2} \square - k$ or $-\frac{4}{3} \square - k$ or $\frac{1}{3} \square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1 : One correct answer $(k < \frac{1}{2}, k \le \frac{1}{3}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ if no method or working (as shown above) then A0 c2M1dep : Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2} \square - k$ and $-\frac{4}{3} \square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1 : Both correct answers only $(k < \frac{1}{2} \text{ or } k + \frac{1}{2} \text{ and } k > \frac{4}{3} \text{ or } k > \frac{4}{3}$ with working as shown above	80	64	100	80								
b i i b i i i i b i i i b i i c i b i i b i i b i i b i i b i i d i d e e e e e d e i e e d e i e f i i f i i i f i i i i i i i i i i	90	72	112.5	90								
b12B1: cao <i>V</i> labelled – dependent on first three B marks in (a) and the first B mark in (b) biilM1: Solving correct pair of simultaneous equations for their <i>V</i> (or if not labelled then the vertex consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective line biilA1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bii2A1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). Objective line method: $-\frac{1}{2} \square - k$ or $-\frac{4}{3} \square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not {e}, \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3}$ or if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2} \square - k$ and $-\frac{4}{3} \square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not f(\frac{1}{2} \text{ or } k \not f(\frac{1}{2} \text{ or } k \not f(\frac{1}{3} \text{ or } k^3 - \frac{4}{3})$ with working as shown above	100	80	125	100								
billM1: Solving correct pair of simultaneous equations for their <i>V</i> (or if not labelled then the vertex consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective line billA1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bill2A1: cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b) SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). Objective line method: $-\frac{1}{2}\square - k$ or $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not {\bf E}, \frac{1}{2}, k > \frac{4}{3}, k^3, \frac{4}{3})$ if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not {\bf E}, \frac{1}{2}$ and $k > \frac{4}{3}$ or $k^3 - \frac{4}{3}$ with working as shown above	bi2BI: ca	o V labelled –	dependent	on first th	ree B marks in (a) and the first B mark in (b)	,						
consistent with their objective line) – this mark can be implied by (42, 44) but in all cases they must have drawn four lines with at least two correct and an attempt at an objective line bii1A1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bii2A1: cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b) SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). <u>Objective line method</u>: $-\frac{1}{2}\square - k$ or $-\frac{4}{3}\square - k$ or $\frac{1}{3}\square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not {n} \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3}$) - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not {n} \frac{1}{2}$ and $k > \frac{4}{3}$ or $k^3 - \frac{4}{3}$) with working as shown above		olving correct	pair of sir	nultaneous	equations for their V (or if not labelled then the	vertex						
bii A1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bii 2A1: cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b) SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \ kx_1 + y_1$ or $60k + 20 \ kx_1 + y_1$ or $42k + 44 \ kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is their numerical V or (42, 44). Objective line method: $-\frac{1}{2}\Box - k$ or $-\frac{4}{3}\Box - k$ or $\frac{1}{3}\Box - k$ where \Box is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not \in \frac{1}{2}, k > \frac{4}{3}, k^3 \ \frac{4}{3})$ - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \ kx_1 + y_1$ and $60k + 20 \ kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their V (but not ((10, 60) or (60, 20)) (so V must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answer stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not \in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$ with working as shown above	drown fou	with their obj	loost two) – tills illa	rk can be implied by (42, 44) but in all cases the	/ must have						
bit A1: cao ((42, 44) only) – dependent on first three B marks in (a) and the first B mark in (b) bit 2A1: cao (77.6 only) – dependent on first three B marks in (a) and the first B mark in (b) SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 kx_1 + y_1 \text{ or } 60k + 20 kx_1 + y_1 \text{ or } 42k + 44 kx_1 + y_1 \text{ where } is any inequality sign or the equals sign and (x_1, y_1) is their numerical V or (42, 44). Objective line method:-\frac{1}{2} - k \text{ or } -\frac{4}{3} - k \text{ or } \frac{1}{3} - k \text{ where } is any inequality or equals. Or one correct answer statedc1A1: One correct answer (k < \frac{1}{2}, k \notin \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3}) if no method or working (as shown above) then A0c2M1dep: Point testing: 10k + 60 kx_1 + y_1 \text{ and } 60k + 20 kx_1 + y_1 \text{ where } is any inequality sign or theequals sign and (x_1, y_1) is (42, 44) or their V (but not ((10, 60) or (60, 20)) (so V must now be theintersection of the two lines 4x + 3y = 300 and x + 2y = 130). Objective line: -\frac{1}{2} - k and -\frac{4}{3} - k where is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only (k < \frac{1}{2} \text{ or } k \pounds \frac{1}{2} \text{ and } k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3} with working as shown above$		((42, 44)) or	least two (ndent on fi	rst three B marks in (a) and the first B mark in (h)						
SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). Objective line method: $-\frac{1}{2}\square - k$ or $-\frac{4}{3}\square - k$ or $\frac{1}{3}\square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \notin \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$ with working as shown above	$bii 2 \Delta 1 \cdot c$	((+2, ++))	_ denende	nt on first i	three B marks in (a) and the first B mark in (b)	J)						
SC in (b) if no objective line drawn then can score in (b) B0B0M1A1A0 for both (42, 44) and 77.6 only provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). Objective line method: $-\frac{1}{2}\square - k$ or $-\frac{4}{3}\square - k$ or $\frac{1}{3}\square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not\in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$) with working as shown above	0112741.00	to (77.0 omy)	ucpende		the D marks in (a) and the first D mark in (b)							
provided that the first three B marks earned in (a) Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \ kx_1 + y_1$ or $60k + 20 \ kx_1 + y_1$ or $42k + 44 \ kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is their numerical V or (42, 44). Objective line method: $-\frac{1}{2}\Box - k \text{ or } -\frac{4}{3}\Box - k \text{ or } \frac{1}{3}\Box - k$ where \Box is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not (\frac{1}{2}, k) / (\frac{4}{3}, k)$	SC in (b)	if no objectiv	e line drav	wn then ca	on score in (b) R0R0M1A1A0 for both (42–44)	and 77.6 only						
Marks in part (c) are not dependent on the marks in (a) and/or (b) c1M1: Point testing method: $10k + 60 \square kx_1 + y_1$ or $60k + 20 \square kx_1 + y_1$ or $42k + 44 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is their numerical V or (42, 44). Objective line method: $-\frac{1}{2}\square - k \text{ or } -\frac{4}{3}\square - k \text{ or } \frac{1}{3}\square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not \in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their V (but not ((10, 60) or (60, 20)) (so V must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not \in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$) with working as shown above	provided	that the first	three B m	arks earn	ed in (a)	und / / to only						
c1M1: Point testing method: $10k + 60 \ kx_1 + y_1$ or $60k + 20 \ kx_1 + y_1$ or $42k + 44 \ kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or $(42, 44)$. Objective line method: $-\frac{1}{2}\Box - k$ or $-\frac{4}{3}\Box - k$ or $\frac{1}{3}\Box - k$ where \Box is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 \ \frac{4}{3})$ - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \ kx_1 + y_1$ and $60k + 20 \ kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is $(42, 44)$ or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not\in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 \ \frac{4}{3}$) with working as shown above	Marks in	part (c) are n	ot depend	lent on the	e marks in (a) and/or (b)							
any inequality sign or the equals sign and (x_1, y_1) is their numerical <i>V</i> or (42, 44). Objective line method: $-\frac{1}{2}\square - k \text{ or } -\frac{4}{3}\square - k \text{ or } \frac{1}{3}\square - k$ where \square is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not\in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$) with working as shown above	c1M1: Po	int testing met	hod: 10k +	$-60 \square kx_1 +$	$v, \text{ or } 60k + 20 \square kx_1 + v, \text{ or } 42k + 44 \square kx_1 + v_1$	where \Box is						
- $\frac{1}{2}$ - k or - $\frac{4}{3}$ - k or $\frac{1}{3}$ - k where \Box is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60\Box kx_1 + y_1$ and $60k + 20\Box kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their V (but not ((10, 60) or (60, 20)) (so V must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not\in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$) with working as shown above	any inequa	ality sign or th	e equals si	gn and (x_1)	v_1 is their numerical V or (42, 44). Objective li	ne method:						
- $\frac{1}{2}$ - k or - $\frac{1}{3}$ - k or $\frac{1}{3}$ - k where \Box is any inequality or equals. Or one correct answer stated c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 - \frac{4}{3})$ - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \Box kx_1 + y_1$ and $60k + 20 \Box kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not\in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$) with working as shown above	1	4	1									
c1A1: One correct answer $(k < \frac{1}{2}, k \not\in \frac{1}{2}, k > \frac{4}{3}, k^3 , \frac{4}{3})$ - if no method or working (as shown above) then A0 c2M1dep: Point testing: $10k + 60 \Box kx_1 + y_1$ and $60k + 20 \Box kx_1 + y_1$ where \Box is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not\in \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 - \frac{4}{3}$) with working as shown above	$-\frac{1}{2}\Box - k$	$-\frac{1}{2}$ - k or $-\frac{4}{3}$ - k or $\frac{1}{3}$ - k where \Box is any inequality or equals. Or one correct answer stated										
c2M1dep: Point testing: $10k + 60 \square kx_1 + y_1$ and $60k + 20 \square kx_1 + y_1$ where \square is any inequality sign or the equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\square - k$ and $-\frac{4}{3}\square - k$ where \square is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \pounds \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 \frac{4}{3}$) with working as shown above	c1A1: One correct answer $(k < \frac{1}{2}, k \notin \frac{1}{2}, k > \frac{4}{3}, k^3 + \frac{4}{3})$ - if no method or working (as shown above) then A0											
equals sign and (x_1, y_1) is (42, 44) or their <i>V</i> (but not ((10, 60) or (60, 20)) (so <i>V</i> must now be the intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \pounds \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 \frac{4}{3}$) with working as shown above	c2M1dep: Point testing: $10k + 60 \square kx_1 + v_1$ and $60k + 20 \square kx_1 + v_1$, where \square is any inequality sign or the											
intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$ where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \pounds \frac{1}{2}$ and $k > \frac{4}{3} \text{ or } k^3 \frac{4}{3}$) with working as shown above	equals sign and (x_1, y_1) is (42, 44) or their V (but not ((10, 60) or (60, 20)) (so V must now be the											
where \Box is any inequality or equals. Or both correct answers stated with no working c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \pounds \frac{1}{2} \text{ and } k > \frac{4}{3} \text{ or } k^3 \frac{4}{3})$ with working as shown above	intersection of the two lines $4x + 3y = 300$ and $x + 2y = 130$). Objective line: $-\frac{1}{2}\Box - k$ and $-\frac{4}{3}\Box - k$											
c2A1: Both correct answers only $(k < \frac{1}{2} \text{ or } k \not \Sigma \frac{1}{2} \text{ and } k > \frac{4}{3} \text{ or } k^3 \frac{4}{3})$ with working as shown above	where \Box is any inequality or equals. Or both correct answers stated with no working											
	c2A1: Bot	th correct answ	vers only ($k < \frac{1}{2}$ or k	$\pm \frac{1}{2}$ and $k \ge \frac{4}{3}$ or $k \ge \frac{4}{3}$) with working as shown a	ibove						

Question Number	Scheme										Mar	ks		
7. (a)	3 < -	$3 < \frac{228}{n} \le 4$										M1		
	Criti	cal va	alue c	of 57	and 7	6 (or	57 ai	nd 75)				A1	
	57≤	n < 7	76 (oi	r 57≤	$\leq n \leq n$	75)			<u> </u>				A1	(3)
	e.g. 1	niddl	le rigl	ht										
	14 20 23 17 15 22 19 25 13 28 32									M1				
	23	25	28	32	<u>22</u>	14	20	17	15	19	13		1411	
	32	<u>28</u>	23	25	<u>22</u>	20	17	19	<u>15</u>	14	13		A1	
	32	<u>28</u>	<u>25</u>	23	<u>22</u>	20	19	<u>17</u>	<u>15</u>	14	<u>13</u>		A1ft	
	32	<u>28</u>	<u>25</u>	23	<u>22</u>	20	<u>19</u>	<u>17</u>	<u>15</u>	14	<u>13</u>		A1	(4)
	e.g. 1	niddl	le left	-										
(b)	14	20	23	17	15	22	19	25	13	28	32			
	23	25	28	32	<u>22</u>	14	20	17	15	19	13			
	28	32	<u>25</u>	23	<u>22</u>	20	19	<u>17</u>	14	15	13			
	32	<u>28</u>	<u>25</u>	23	<u>22</u>	<u>20</u>	19	<u>17</u>	<u>15</u>	14	13			
	32	<u>28</u>	<u>25</u>	23	<u>22</u>	<u>20</u>	19	<u>17</u>	<u>15</u>	<u>14</u>	13			
(c)	(c) From first-fit Bin 1 could not fit the 17 so $n < 74$ (or $n \le 73$) but could fit the 15 so <i>n</i> is either 72 (as the largest total is 72 in Bin 1 from first-fit) or 73						73) but could fit the 15 rst-fit) or 73	B1						
	Fron	n first	t-fit d	ecrea	sing	the 13	3 cou	ld no	t fit ir	n Bin	1		B1	
	So n	= 72											ddB1	(3)
													10 mar	ks

Question	Scheme	Marks				
Number						
	Notes for Question 7					
a1M1: Ar	equation or inequality linking the expression $\frac{228}{n}$ with either 3 or 4					
a1A1: Co	rrect critical values of 57 and 76 (or 57 and 75)					
a2A1: 5 7	$\leq n < 76 \text{ or } 57 \leq n \leq 75$					
b1M1: Qu M1 only. 1 b1A1: Fir b2A1ft: S b3A1: csc	tick sort – pivots, p, selected and first pass gives >p, p, <p. 1="" choosing="" if="" only="" pivot<br="">If sorting into ascending order then mark as a misread st pass correct and next pivots chosen correctly/consistently for second pass econd and third passes correct (ft from their first pass and choice of pivots) o (including a fourth pass with 19 used as a pivot if middle right or 14 if middle left)</p.>	per iteration				
c1B1: Con may see (1 realising t n^3 72	c1B1: Correct deduction from first-fit that <i>n</i> is at least 72 or at most 73 (oe e.g. less than 74). For example, may see $(14 + 20 + 23 + 15 =)$ 72 stated therefore n^3 72 or $14 + 20 + 23 + 17 = 74$ followed by $n < 74$ (so realising that the 17 did not fit in Bin 1). As a minimum accept the statement that $n < 74$ or $n \le 73$ or n^3 72					
c2B1: Cor 32 + 28 + or simply c3ddB1: of they need so <i>n</i> cannot talking ab	rect deduction from first-fit decreasing that the 13 was not placed in Bin 1. For example 13 = 73 so therefore $n < 73$ or $n \pounds$ 72. As a minimum accept the statement that $n <$ stating that '13 did not fit in Bin 1' (give bod here if not clear which Bin 1 they are cao (dependent on both previous B marks) – must state that the <u>largest</u> total in any b to say or show there exists a bin with 72) and that the <u>13 did not fit in Bin 1 in first</u> of be <u>73</u> and therefore $n = 72$ (not just '13 does not fit in Bin 1' – must be clear that out first-fit decreasing)	mple, may see 73 or <i>n</i> £ 72 considering) bin is <u>72</u> (or <u>t-fit decreasing</u> they are				
No marks	in (c) if $n = 72$ stated with no working or if all the candidate does is to sum the num	obers in each				

No marks in (c) if n = 72 stated with no working or if all the candidate does is to sum the numbers in each bin

Note that the first B mark in (c) can be implied if the candidate considers the first-fit decreasing packing first or argues with first-fit decreasing before considering first-fit, e.g., 'The 13 does not fit in Bin 1 in the first-fit decreasing packing therefore n is at most 72 and the total of Bin 1 in first-fit is 72' would imply the first two B marks in this part. Stating that therefore n must be 72 would then score all three marks

Pearson Education Limited. Registered company number 872828 with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom